



For people with hard-to-treat HIV

First-of-its-kind
SUNLENCA[®]
is a **twice-**
a-year,
injectable
treatment
used in combination
with other HIV-1
medicines

THIS COULD BE YOUR SHOT

Ask your healthcare provider
if SUNLENCA is right for you.

What is SUNLENCA?

SUNLENCA is a prescription medicine that is used with other human immunodeficiency virus-1 (HIV-1) medicines to treat HIV-1 infection in adults:

- who have received HIV-1 medicines in the past, **and**
- who have HIV-1 virus that is resistant to many HIV-1 medicines, **and**
- whose current HIV-1 medicines are failing. Your HIV-1 medicines may be failing because the HIV-1 medicines are not working or no longer work, you are not able to tolerate the side effects, or there are safety reasons why you cannot take them.

SUNLENCA does not cure HIV-1 or AIDS. HIV-1 is the virus that causes AIDS.

IMPORTANT SAFETY INFORMATION

Who should not receive or take SUNLENCA?

Do not receive or take SUNLENCA if you take other medicines called strong CYP3A inducers. Ask your healthcare provider if you are not sure.

Please click for [Important Facts about SUNLENCA](#).

 Sunlenca[®]
(lenacapavir) injection
463.5 mg/1.5 mL



HIV treatments continue to evolve




If you are living with hard-to-treat HIV, you know how frustrating it can feel when your HIV treatments stop working. Now, there's another option available that may be able to help treat your HIV.

If your provider decides SUNLENCA is right for you, they may also adjust the combination of other HIV medicines you take.

What is SUNLENCA?

SUNLENCA is an injectable treatment for HIV with a twice-a-year dosing schedule. SUNLENCA can help you treat your HIV when used in combination with the other HIV-1 medicines that have been prescribed to you by your doctor.

Adding SUNLENCA may be right for you if:

-  you have received HIV-1 medicines in the past, **and**
-  you have HIV-1 virus that is resistant to many HIV-1 medicines, **and**
-  your current HIV-1 medicines are failing. Your HIV-1 medicines may be failing because the HIV-1 medicines are not working or no longer work, you are not able to tolerate the side effects, or there are safety reasons why you cannot take them.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of SUNLENCA?

Serious side effects of SUNLENCA may include:

- **Changes in your immune system.** Your immune system may get stronger and begin to fight infections that may have been hidden in your body. Tell your healthcare provider if you have any new symptoms after you start taking SUNLENCA.
- **Injection site reactions** may happen and may include swelling, pain, redness, skin hardening, small mass or lump, and itching. Hardened skin or lumps at the injection site usually can be felt but not seen. If you develop hardened skin or a lump, it may take longer than other reactions at the injection site to go away, and the injection site may not completely heal on its own. Tell your healthcare provider if you have any injection site reactions.

The most common side effects of SUNLENCA were nausea and injection site reactions. These are not all the possible side effects. Tell your healthcare provider if you have any side effects that bother you or don't go away.

HIV treatments continue to evolve

If you are living with hard-to-treat HIV, you know how frustrating it can feel when your HIV treatments stop working. Now, there's another option available that may be able to help treat your HIV.

If your provider decides SUNLENCA is right for you, they may also adjust the combination of other HIV medicines you take.

Between your 6-month injections, SUNLENCA keeps working in your body



SUNLENCA is a subcutaneous injection, meaning it's injected just under the skin and does not go deeply into the muscle. If you are prescribed SUNLENCA, the injections will always be given to you by a healthcare provider and may even be given at your regular appointments you may have already scheduled.



SUNLENCA is long-acting. The medicine will continue to work in your body during the 6 months between your maintenance appointments.* Make sure to attend your appointments as scheduled.

- ● ● **Remember, SUNLENCA works with your other HIV-1 medications. Continue to take all of your other HIV-1 medicines as prescribed by your doctor.**

*Maintenance appointments can be scheduled up to 2 weeks before or after the 6-month mark. If you miss a SUNLENCA appointment, contact your healthcare provider as soon as possible. They'll determine if and how you should continue with SUNLENCA.

IMPORTANT SAFETY INFORMATION (cont'd)

What should I tell my healthcare provider before taking SUNLENCA?

- **All your health problems.**
- **All the medicines you take**, including prescription and over-the-counter medicines, antacids, laxatives, vitamins, and herbal supplements, including St. John's wort. SUNLENCA and other medicines may affect each other. Keep a list of all your medicines and show it to your healthcare provider and pharmacist, and ask if it is safe to take SUNLENCA with all of your other medicines. Do not start other medicines without telling your healthcare provider. SUNLENCA may affect certain other medicines for up to 9 months after your last injection.
- **If you are pregnant** or plan to become pregnant. It is not known if SUNLENCA can harm your unborn baby. Tell your healthcare provider if you become pregnant while taking SUNLENCA.
- **If you are breastfeeding** (nursing) or plan to breastfeed. Do not breastfeed. HIV-1 can be passed to the baby in breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.




Getting started with SUNLENCA


There are two approaches to starting treatment with SUNLENCA. Your healthcare provider will discuss the options for starting SUNLENCA with you.

To start SUNLENCA, you will take SUNLENCA pills (300 mg each) in addition to receiving injections. SUNLENCA pills will be delivered through a specialty pharmacy to either your home or your provider's office. Be sure to work with your healthcare team to coordinate delivery from the specialty pharmacy.

Regardless of which initiation option you use (as shown on the right), see the next page for how to continue taking SUNLENCA




Initiation Option 1: Same-day start



● ● ● Remember, SUNLENCA works with your other HIV-1 medications. Continue to take all of your other HIV-1 medicines as prescribed by your doctor.

Initiation Option 2: Step-by-step start



● ● ● Remember, SUNLENCA works with your other HIV-1 medications. Continue to take all of your other HIV-1 medicines as prescribed by your doctor.

IMPORTANT SAFETY INFORMATION

Who should not receive or take SUNLENCA?

Do not receive or take SUNLENCA if you take other medicines called strong CYP3A inducers. Ask your healthcare provider if you are not sure.




Getting started with SUNLENCA

There are two approaches to starting treatment with SUNLENCA. Your healthcare provider will discuss the options for starting SUNLENCA with you.

To start SUNLENCA, you will take SUNLENCA pills (300 mg each) in addition to receiving injections. SUNLENCA pills will be delivered through a specialty pharmacy to either your home or your provider's office. Be sure to work with your healthcare team to coordinate delivery from the specialty pharmacy.

Maintenance Schedule: Continuing treatment

Once you've started SUNLENCA, you will get 2 injections from your healthcare provider every 6 months.* Make sure to attend your appointments as scheduled.



- ● ● **Remember, SUNLENCA works with your other HIV-1 medications. Continue to take all of your other HIV-1 medicines as prescribed by your doctor.**

*Maintenance appointments can be scheduled up to 2 weeks before or after the 6-month mark. If you miss a SUNLENCA appointment, contact your healthcare provider as soon as possible. They'll determine if and how you should continue with SUNLENCA.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of SUNLENCA?

Serious side effects of SUNLENCA may include:



- **Changes in your immune system.** Your immune system may get stronger and begin to fight infections that may have been hidden in your body. Tell your healthcare provider if you have any new symptoms after you start taking SUNLENCA.
- **Injection site reactions** may happen and may include swelling, pain, redness, skin hardening, small mass or lump, and itching. Hardened skin or lumps at the injection site usually can be felt but not seen. If you develop hardened skin or a lump, it may take longer than other reactions at the injection site to go away, and the injection site may not completely heal on its own. Tell your healthcare provider if you have any injection site reactions.

The most common side effects of SUNLENCA were nausea and injection site reactions. These are not all the possible side effects. Tell your healthcare provider if you have any side effects that bother you or don't go away.

TRACK YOUR APPOINTMENTS

If you and your healthcare provider have chosen the same-day start option, use this chart to help track your start with SUNLENCA[®].

MY START SCHEDULE: OPTION 1

<div style="background-color: #003366; color: white; padding: 5px; border-radius: 10px; margin-bottom: 5px;">Day 1</div>  <p>DATE: _____</p>	<div style="background-color: #003366; color: white; padding: 5px; border-radius: 10px; margin-bottom: 5px;">Day 2</div>  <p>DATE: _____</p>	<p>During the 6 months between injections, be sure to continue taking your other HIV-1 medications.</p>	<div style="background-color: #003366; color: white; padding: 5px; border-radius: 10px; margin-bottom: 5px;">Maintenance appointment</div> <p>DATE: _____</p> <p>You will begin maintenance doses 6 months after your first set of injections.*</p>
--	--	--	---

Continue to take other HIV-1 medications as prescribed by your healthcare provider.

<ul style="list-style-type: none"> • Take 2 SUNLENCA pills (300 mg each) • Receive 2 SUNLENCA injections 	<ul style="list-style-type: none"> • Take 2 SUNLENCA pills 	<ul style="list-style-type: none"> • Receive 2 SUNLENCA injections
--	---	---

*Maintenance appointments can be scheduled up to 2 weeks before or after the 6-month mark. If you miss a SUNLENCA appointment, contact your healthcare provider as soon as possible. They'll determine if and how you should continue with SUNLENCA.



USE THIS SPACE TO WRITE DOWN THE OTHER MEDICATIONS YOU WILL TAKE IN COMBINATION WITH SUNLENCA:

IMPORTANT SAFETY INFORMATION (cont'd)

What should I tell my healthcare provider before taking SUNLENCA?





- **All your health problems.**
- **All the medicines you take**, including prescription and over-the-counter medicines, antacids, laxatives, vitamins, and herbal supplements, including St. John's wort. SUNLENCA and other medicines may affect each other. Keep a list of all your medicines and show it to your healthcare provider and pharmacist, and ask if it is safe to take SUNLENCA with all of your other medicines. Do not start other medicines without telling your healthcare provider. SUNLENCA may affect certain other medicines for up to 9 months after your last injection.
- **If you are pregnant** or plan to become pregnant. It is not known if SUNLENCA can harm your unborn baby. Tell your healthcare provider if you become pregnant while taking SUNLENCA.
- **If you are breastfeeding** (nursing) or plan to breastfeed. Do not breastfeed. HIV-1 can be passed to the baby in breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

TRACK YOUR APPOINTMENTS

If you and your healthcare provider have chosen the step-by-step start option, use this chart to help track your start with SUNLENCA[®].

MY START SCHEDULE: OPTION 2

<div style="background-color: #002060; color: white; padding: 5px; border-radius: 10px; margin-bottom: 5px;">Day 1</div>  <p>DATE: _____</p>	<div style="background-color: #002060; color: white; padding: 5px; border-radius: 10px; margin-bottom: 5px;">Day 2</div>  <p>DATE: _____</p>	<div style="background-color: #002060; color: white; padding: 5px; border-radius: 10px; margin-bottom: 5px;">Day 8</div>  <p>DATE: _____</p>	<div style="background-color: #002060; color: white; padding: 5px; border-radius: 10px; margin-bottom: 5px;">Day 15</div>  <p>DATE: _____</p>	<p>During the 6 months between injections, be sure to continue taking your other HIV-1 medications.</p>
--	--	--	---	--

Continue to take other HIV-1 medications as prescribed by your healthcare provider.

<ul style="list-style-type: none"> • Take 2 SUNLENCA pills (300 mg each) 	<ul style="list-style-type: none"> • Take 2 SUNLENCA pills 	<ul style="list-style-type: none"> • Take 1 SUNLENCA pill 	<ul style="list-style-type: none"> • Receive 2 SUNLENCA injections
---	---	--	---

Maintenance appointment

DATE: _____

You will begin maintenance doses 6 months after your first set of injections.*

- Receive 2 SUNLENCA injections

*Maintenance appointments can be scheduled up to 2 weeks before or after the 6-month mark. If you miss a SUNLENCA appointment, contact your healthcare provider as soon as possible. They'll determine if and how you should continue with SUNLENCA.



USE THIS SPACE TO WRITE DOWN THE OTHER MEDICATIONS YOU WILL TAKE IN COMBINATION WITH SUNLENCA:

IMPORTANT SAFETY INFORMATION (cont'd)

Who should not receive or take SUNLENCA?

Do not receive or take SUNLENCA if you take other medicines called strong CYP3A inducers. Ask your healthcare provider if you are not sure.



If your healthcare provider has already determined that a Gilead treatment is right for you

The Gilead Advancing Access[®] program is committed to helping you afford your medication, no matter your situation.

Whether you are insured or uninsured, Advancing Access may be able to help in the following ways:



If eligible, Advancing Access may be able to help you lower your co-pay to as little as \$0*



Identify potential sources of financial support if you have government insurance or are uninsured



For more information, visit [GileadAdvancingAccess.com](https://www.gileadadvancingaccess.com) or call **1-800-226-2056**, M-F, 9 AM to 8 PM ET

*Eligible, commercially insured individuals enrolled in Gilead Advancing Access co-pay coupon card program could pay as little as \$0. Restrictions apply. Subject to change. See full terms and conditions at [GileadAdvancingAccess.com](https://www.gileadadvancingaccess.com).

● ● ● **Share your story. Inspire others. Become an advocate. If interested, please call us at 1-888-235-7070 or send us an email at info@GANteam.com to get started.**



SUNLENCA[®] may be your shot

Help set yourself up for success with this list of helpful reminders that outline the next steps after being prescribed SUNLENCA.

A specialty pharmacy is responsible for shipping SUNLENCA. Your healthcare team will work with you to coordinate the details.

- 1 Remember to continue taking your other HIV-1 medications during SUNLENCA treatment, including during the periods of time between injections. Do not miss a dose. You should always take the HIV medications in your treatment regimen exactly as instructed by your healthcare provider, including SUNLENCA
- 2 Establish your start plan when your healthcare provider prescribes SUNLENCA to ensure a smooth transition onto treatment. If necessary, set up the specialty pharmacy details at the same appointment. The specialty pharmacy may be different than your usual pharmacy
- 3 After you establish your start plan with your healthcare provider, fill out the treatment tracker for **Initiation Option 1** or **Initiation Option 2** to help you stay on track with SUNLENCA. Make sure you follow your schedule and attend all of your appointments

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of SUNLENCA?

Serious side effects of SUNLENCA may include:

- **Changes in your immune system.** Your immune system may get stronger and begin to fight infections that may have been hidden in your body. Tell your healthcare provider if you have any new symptoms after you start taking SUNLENCA.
- **Injection site reactions** may happen and may include swelling, pain, redness, skin hardening, small mass or lump, and itching. Hardened skin or lumps at the injection site usually can be felt but not seen. If you develop hardened skin or a lump, it may take longer than other reactions at the injection site to go away, and the injection site may not completely heal on its own. Tell your healthcare provider if you have any injection site reactions.

The most common side effects of SUNLENCA were nausea and injection site reactions. These are not all the possible side effects. Tell your healthcare provider if you have any side effects that bother you or don't go away.

THIS COULD BE YOUR SHOT

Visit [SUNLENCA.com](https://www.sunlenca.com) to learn more

IMPORTANT SAFETY INFORMATION (cont'd)

What should I tell my healthcare provider before taking SUNLENCA?

- **All your health problems.**
- **All the medicines you take**, including prescription and over-the-counter medicines, antacids, laxatives, vitamins, and herbal supplements, including St. John's wort. SUNLENCA and other medicines may affect each other. Keep a list of all your medicines and show it to your healthcare provider and pharmacist, and ask if it is safe to take SUNLENCA with all of your other medicines. Do not start other medicines without telling your healthcare provider. SUNLENCA may affect certain other medicines for up to 9 months after your last injection.
- **If you are pregnant** or plan to become pregnant. It is not known if SUNLENCA can harm your unborn baby. Tell your healthcare provider if you become pregnant while taking SUNLENCA.
- **If you are breastfeeding** (nursing) or plan to breastfeed. Do not breastfeed. HIV-1 can be passed to the baby in breast milk.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please click for Important Facts about [SUNLENCA](#).



SUNLENCA, the SUNLENCA Logo, ADVANCING ACCESS, GILEAD, and the GILEAD Logo are trademarks of Gilead Sciences, Inc., or its related companies.

© 2023 Gilead Sciences, Inc. All rights reserved. US-LENC-0023 01/23

(sun-LEN-kuh)

IMPORTANT FACTS

This is only a brief summary of important information about SUNLENCA® and does not replace talking to your healthcare provider about your condition and your treatment.

ABOUT SUNLENCA

SUNLENCA is a prescription medicine that is used with other human immunodeficiency virus-1 (HIV-1) medicines to treat HIV-1 infection in adults:

- who have received HIV-1 medicines in the past, **and**
- who have HIV-1 virus that is resistant to many HIV-1 medicines, **and**
- whose current HIV-1 medicines are failing. Your HIV-1 medicines may be failing because the HIV-1 medicines are not working or no longer work, you are not able to tolerate the side effects, or there are safety reasons why you cannot take them.

SUNLENCA does not cure HIV-1 or AIDS. HIV-1 is the virus that causes AIDS.

Do NOT receive or take SUNLENCA if you also take certain other medicines called strong CYP3A inducers. Ask your healthcare provider if you are not sure.

BEFORE RECEIVING OR TAKING SUNLENCA

Tell your healthcare provider all your medical conditions, including if you:

- Are pregnant or plan to become pregnant.
- Are breastfeeding (nursing) or plan to breastfeed. Do not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, including St. John's wort, and show it to your healthcare provider and pharmacist.
- Ask your healthcare provider or pharmacist about medicines that interact with SUNLENCA.
- SUNLENCA may affect certain other medicines for up to 9 months after your last injection.

POSSIBLE SIDE EFFECTS OF SUNLENCA

SUNLENCA can cause serious side effects, including:

- Changes in your immune system.
- Injection site reactions, including swelling, pain, redness, skin hardening, small mass or lump, and itching. Hardened skin or lumps at the injection site usually can be felt but not seen. Tell your healthcare provider if you have any injection site reactions.
- **The most common side effects of SUNLENCA** are nausea and injection site reactions.

These are not all the possible side effects of SUNLENCA. Tell your healthcare provider right away if you have any new symptoms while taking SUNLENCA.

HOW TO RECEIVE AND TAKE SUNLENCA

SUNLENCA treatment will consist of injections and tablets. SUNLENCA injections will be given to you by your healthcare provider under the skin (subcutaneous injection) in your stomach-area (abdomen). Take SUNLENCA tablets by mouth, with or without food.

There are two options (Option 1 and Option 2) to start treatment with SUNLENCA. Your healthcare provider will decide which starting option is for you.

- If Option 1 is chosen:
 - On Day 1, you will receive 2 SUNLENCA injections and take 2 SUNLENCA tablets.
 - On Day 2, you will take 2 SUNLENCA tablets.
- If Option 2 is chosen:
 - On Day 1 and Day 2, you will take 2 SUNLENCA tablets each day.
 - On Day 8, you will take 1 SUNLENCA tablet.
 - On Day 15, you will receive 2 SUNLENCA injections.

After completing Option 1 or Option 2, you will receive 2 SUNLENCA injections every 6 months (26 weeks) from the date of your last injection. If you miss your scheduled injection appointment, call your healthcare provider right away to discuss your treatment options.

GET MORE INFORMATION

- This is only a brief summary of important information about SUNLENCA. Talk to your healthcare provider or pharmacist to learn more.
- Go to [SUNLENCA.com](https://www.sunlenca.com) or call 1-800-GILEAD-5.
- If you need help paying for your medicine, visit [SUNLENCA.com](https://www.sunlenca.com) for program information.